

Psychiatric Disorders and Psychotherapy of Substance Abuse

Robert M. Weinrieb, M.D
Department of Psychiatry
University of Pennsylvania

Pre-Lecture Exam

Question 1

- 1. Which anxiety symptom is most commonly associated with primary alcoholism?**
 - a. Panic while drinking**
 - b. Panic while sober**
 - c. General Anxiety Disorder while drinking**
 - d. Withdrawal palpitations and/or shortness of breath**
 - e. Agoraphobia while intoxicated or in withdrawal**

Question 2

- 2. Which of the following is the most common comorbid condition with substance abuse disorders?**
- a. Antisocial Personality Disorder**
 - b. Bipolar Disorder**
 - c. Generalized Anxiety Disorder**
 - d. Agoraphobia**
 - e. Dementia**

Question 3

True or False

People who present for treatment for a substance use disorder are two times more likely to have a second psychiatric disorder than those without a SUD.

Question 4

True or False

Substance use disorder reduces life expectancy by 5 years.

Question 5

True or False

Sixty percent of individuals treated for cocaine dependence are clean six months after acute treatment.

Question 6

True or False

Psychotherapy in the context of therapeutic communities has been found to be effective for opiate use disorder.

Attitudes Toward the Treatment of Addicts

At completion of residency, more physicians have negative attitudes toward SUD pts and are less optimistic about benefits of treatment than at the start of med school

--Geller, et al, 1989

So, Why is That ?

- 1. Historically, substance abuse disorders (SUDs) were treated independently of medical community by paraprofessionals**
- 2. Mental health services also rejected pts with SUDs**
- 3. House staff see recidivist patients with multiple complex problems and are not trained to deal with them**

“Attitude Adjustment”

- 1. An adequate knowledge base**
- 2. A positive attitude toward the patient and the benefits of treatment**
- 3. A sense of responsibility for the clinical problem**

- J. A. Renner, Jr. *Biol Psychiatry*, 2004

Topics to be Discussed

1. Dual Diagnosis

- Definition, epidemiology, a case
- Effects on medical care outcomes

2. Psychotherapy of Addiction

- Theory, examples, outcomes

Definition of Dual Diagnosis

- **Dual Diagnosis is defined by having a major psychiatric diagnosis comorbid with a Substance Use Disorder (SUD)**
- **Psychiatric symptoms are common in the context of substance abuse**
- **2/3 individuals with SUD have another psychiatric syndrome (Axis I)**

Dual Diagnosis Caveats

- **Many of these psychiatric syndromes are temporary**

Psychiatric Symptoms Due to Acute Effects of Drugs, ETOH

- Stimulants (cocaine, amphetamines)

Anxiety (panic, PTSD) mania, paranoia, hallucinations, delusions

- Sedative/hypnotics (Etoh, benzos, opiates)

Depression

Psychiatric Symptoms Due to Withdrawal from Drugs, ETOH

- **Stimulants (cocaine, amphetamines)**

Depression


- **Sedative/hypnotics (Etoh, benzos, opiates)**

Anxiety, panic, depression, hallucinosis

Hamilton Depression Score ≥ 20 *

*** For Alcohol**

Source: Brown S, Schuckit M. (1988) J Stud Alcohol;49:412-417.



Anxiety In 171 Primary Alcoholics

Symptom

- Withdrawal palpitations and/or shortness of breath 80%
- Panic while drinking 4%
- Panic while sober 2%
- Generalized anxiety while sober 4%

Source: Brown S, Schuckit M. *J Stud Alcohol.* 1990;51:34-41.

Psychiatric Symptoms: Primary vs. Secondary

- Primary or “Self-Medication Hypotheses”

Independent psychiatric disorder precedes SUD

- Secondary or “The Disease Concept”

Substance induced psychiatric symptoms

- Both are true, but secondary symptoms are more commonly true

Epidemiology of Dual Diagnosis Disorders

- **Epidemiologic Catchment Area Study (ECA)**
- **People who present for treatment for a SUD are ~3 X more likely to have a second psychiatric disorder vs. those without SUD**
- **Most comorbidity (dual diagnosis) is accounted for by Antisocial Personality Disorder (Axis II) and another SUD**

How to Make a Diagnosis When 2+ Disorders Are Observed

1. Take a good history
2. Be able to differentiate among acute and withdrawal symptoms of alcohol and drugs
3. Were psychiatric symptoms present during a clean period of more than 4 weeks?

Case Example of Dual Diagnosis

45 y/o male using cocaine for 5 years. Is depressed with paranoid thoughts. He stabbed himself while trying to fend off an “intruder” in his truck. Brought in by police who witnessed the stabbing-no intruder was seen

Important questions:

1. Did the psychiatric symptoms precede his alcohol dependence?
2. Were there periods of time lasting more than 4 weeks during which psychiatric symptoms were present?
3. Presumptive diagnosis?

Treatment for Dual Diagnosis

- **Integration of therapy is necessary (medications, groups and individual tx)**
- **Sometimes “coercion” or drug courts**
- **Clinical Trials: Seeking Safety (Post-Traumatic Stress D/O)**

Why Improve Medical Care in Dual Diagnosis Patients?

- **SUDs reduce life expectancy by ~14 years**
- **Studies of on-site tx of patients with serious mental illness and SUD found;**
 - 1. Reduced mortality (by up to 1/3)**
 - 2. Increase abstinence from drugs/alcohol**
 - 3. Modest cost**

Initiation of Treatment of SUDs

1. Engagement (Stages of Change)

-Prochaska and DiClemente

- **Precontemplation**
 - **Contemplation**
 - **Action**
 - **Maintenance**

Approach to Treatment of SUDs

- 1. Detoxification**
- 2. Relapse prevention**
- 3. Maintenance of recovery**